arXiv:1304.5018v2 [physics.optics] 24 Apr 2013

Incompleteness of Spin and Orbital Angular Momentum Separation for Light Beams

Marco Ornigotti[1] and Andrea Aiello[1,2*]
[1] *Max Planck Institute for the Science of Light,*
Günther-Scharowsky-Strasse 1/Bau24, 91058 Erlangen, Germany and
[2]*Institute for Optics, Information and Photonics,*
University of Erlangen-Nuernberg, Staudtstrasse 7/B2, 91058 Erlangen, Germany
(Dated: April 25, 2013)

The angular momentum of a circularly polarized beam of light in vacuum is calculated and compared with the value of the same observable for a wave packet of light. While the packet has a finite transverse and longitudinal extent, the span of the beam along the direction of propagation is, in principle, infinite. This simple but subtle fact renders incomplete the textbook calculation of the separation in a spin part and in an orbital part of the angular momentum of a beam of light. We demonstrate that an extra spin-like part must be added in order to preserve the intuitive fact that the spin angular momentum of a circularly polarized beam of light containing N photons cannot exceed the value $\pm N$ times the reduced Planck constant. Finally, the impact of this extra term is quantified by means of two examples: a Laguerre-Gaussian and a Bessel beam, both circularly polarized.

Introduction.—In his *Treatise of Electricity and Magnetism* [1], Maxwell pointed out that the electromagnetic field carries both energy and momentum, and that the momentum can have both linear and angular contributions. Today it is common knowledge that the angular momentum (AM) of light can be thought of having two contributions: a spin part associated to polarization (firstly theoretically investigated by Poynting [2] and then experimentally demonstrated by Beth [3]), and an orbital part associated with the spatial distribution of the field, as first recognised by Darwin [4]. About sixty years later, thanks to the seminal work of Allen and Woerdman [5], the topic of AM of light experienced a new renaissance, producing a considerable amount of literature on the subject, including theoretical discussions about the separation of AM in spin and orbital parts for optical beams [6–14], the spin-Hall effect of light [15–18], the geometric spin-Hall effect of light [19, 20], and a huge variety of experimental applications such as optical tweezers, spatial light modulators and vortex beams. A satisfactory review of the development of the field of optical AM of light is reported in Refs. [21–23].

In this Letter we report on a novel separation of the total AM of a light beam into three parts: the orbital part, the spin part and a new spin-like part which arises from an incomplete two-dimensional integration (over a plane orthogonal to the direction of propagation of the beam) of a three-divergence term. The latter part, also called "surface term" [24], is usually neglected when the light field has a finite extent along any direction. This is not the case for a beam, which is unbounded along the axis of propagation. We refer to this additional term as the *surface spin angular momentum* (SSAM). We show that SSAM arises whenever the intensity of a beam is recorded by a planar detector which, in practice, performs a two dimensional integration over a cross section of the beam perpendicular to its direction of propagation. As this is the case occurring in the majority of experimental optical setups, we believe that the impact of our new results may be significant.

The structure of this Letter is as follows. We first calculate the total AM density $\mathbf{j}(\mathbf{r})$ of an arbitrary electromagnetic wave field. Then, by means of some straightforward manipulations, we decompose $\mathbf{j}(\mathbf{r})$ into the sum of three terms: the well-known orbital and spin AM parts and a third term (the SSAM) which amounts to a three-divergence. Next, we integrate $\mathbf{j}(\mathbf{r})$ upon a two-dimensional surface at $z = const.$, mimicking the detection process, thus obtaining the total AM per unit length. This unravels the nature of the SSAM term and how it manifests in the actual detection of the AM of a light beam. Finally, we calculate the explicit form of the SSAM for the cases of Laguerre-Gaussian and Bessel beams.

Theory of SSAM.—To begin with, by following van Enk and Nienhuis [8] we write the cycle-averaged linear and AM vector densities in vacuo of a monochromatic field of angular frequency ω_0, as

$$\mathbf{p}(\mathbf{r}) = \frac{\varepsilon_0}{2}\mathrm{Re}\left(\mathbf{E}^* \times \mathbf{B}\right), \; \mathbf{j}(\mathbf{r}) = \frac{\varepsilon_0}{2}\mathbf{r} \times \mathrm{Re}\left(\mathbf{E}^* \times \mathbf{B}\right), \quad (1)$$

where the transverse electric field $\mathbf{E}$ and the magnetic field $\mathbf{B}$ are expressible via the gauge invariant transverse part $\mathbf{A}$ of the vector potential as

$$\mathbf{E} = i\omega_0\mathbf{A} \qquad \text{and} \qquad \mathbf{B} = \boldsymbol{\nabla} \times \mathbf{A}, \quad (2)$$

respectively. The complex-valued time-independent fields $\mathbf{A}$, $\mathbf{E}$ and $\mathbf{B}$, are functions of the position vector $\mathbf{r}$ solely. The corresponding time-dependent real-valued fields are given by $\boldsymbol{\mathcal{A}}(\mathbf{r},t) = \mathrm{Re}(\mathbf{A}e^{-i\omega_0 t})$, et cetera. The total linear and angular momenta of the electromagnetic field are given by the integrals of the corresponding densities over all space:

$$\boldsymbol{\mathcal{P}} = \int \mathbf{p}(\mathbf{r})\,\mathrm{d}^3r, \qquad \boldsymbol{\mathcal{J}} = \int \mathbf{j}(\mathbf{r})\,\mathrm{d}^3r. \quad (3)$$

By using Eqs. (1-2) and choosing, throughout this Letter, the physical units in such a way that $\varepsilon_0\omega_0/2 = 1$, we can express the AM density as

$$\begin{aligned}\mathbf{j}(\mathbf{r}) = & - \mathrm{Re}\Big\{ i\mathbf{A}^* \cdot \big(\mathbf{r} \times \boldsymbol{\nabla}\big)\mathbf{A} + \mathbf{A}^* \times \mathbf{A} \\ & - \partial_j \left[A_j^*(\mathbf{r} \times \mathbf{A})\right] \Big\}, \end{aligned} \tag{4}$$

where, according to [24], the first term represents the orbital part of the AM and the second term is the intrinsic or spin AM of light. As it is also noticed in Ref. [24], the third term is a three-divergence that vanishes when integrated over a volume whose surface boundary is so far from the field sources such that $\mathbf{A}(\mathbf{r}) \approx \mathbf{0}$. Thus, if we substitute Eq. (4) in the rightmost expression for $\boldsymbol{\mathcal{J}}$ in Eq. (3), we obtain the textbook expression for the orbital and spin AM separation. However, as pointed out by Haus [25], for a monochromatic beam of light propagating in the z direction the classical optics analogous of the helicity of a photon, namely the projection of the spin AM along the direction of the linear momentum, is given by $\mathcal{S}_z = \hat{\mathbf{z}} \cdot \overline{\mathbf{J}}$ where, henceforth, the over line denotes normalization according to the rule

$$\overline{\mathbf{J}} = \frac{\mathbf{J}}{U/\omega_0}, \tag{5}$$

and with $\mathbf{J}$ and U we have denoted the cycle averaged AM and energy *per unit length*:

$$\mathbf{J} = \int \mathbf{j}(\mathbf{r})\, \mathrm{d}^2R, \qquad U = \omega_0 \int \left|\mathbf{A}(\mathbf{r})\right|^2 \mathrm{d}^2R. \tag{6}$$

In the expressions above $\mathrm{d}^2R = \mathrm{d}x\,\mathrm{d}y$ and the integration is over all the plane xy at $z = const.$ This definition is also consistent with the physical picture of a beam of light as observed on a laboratory bench, where it travels across plates, lenses and other optical devices before eventually shining the surface of a *planar* detector, as opposed to a 3D particle detector.

As a consequence of the surface, as opposed to volume integration, a beam-like field "feels" the contribution given by the z-part of the latter term in Eq. (4), namely $\partial_z[A_z^*(\mathbf{r} \times \mathbf{A})]$ because such term is *not* integrated when one calculates the AM per unit length, namely the integral of the AM density $\mathbf{j}(\mathbf{r})$ over all the xy plane at $z = const.$ In practice, since $\int \partial_j[A_j^*(\mathbf{r} \times \mathbf{A})]\mathrm{d}^2R = \partial_z \int A_z^*(\mathbf{r} \times \mathbf{A})\mathrm{d}^2R$, we can write the SSAM contribution in the following form:

$$\mathbf{J}^{\mathrm{surf}} = \frac{\partial}{\partial z} \int \mathrm{Re}\Big\{ iA_z^*(\mathbf{r} \times \mathbf{A}) \Big\} \mathrm{d}^2R. \tag{7}$$

From Eq. (7) it seems that the $\mathbf{J}^{\mathrm{surf}}$ depends both on the potential vector $\mathbf{A}$ and the position vector $\mathbf{r}$, thus revealing a seemingly hybrid orbital/spin nature. However, in the remainder we will show results that do not support this interpretation.

Now, in order to give the SSAM an explicit form, we use the angular spectrum representation for the vector potential using the so-called helicity basis [26] $\hat{\mathbf{h}}_\sigma(\mathbf{K}) = (1/\sqrt{2})[\hat{\mathbf{e}}_1(\mathbf{K}) + i\sigma\hat{\mathbf{e}}_2^*(\mathbf{K})]$, where $\hat{\mathbf{e}}_{1,2}(\mathbf{K})$ are two unit vectors orthogonal to $\mathbf{k} = \mathbf{K} + k_z\hat{\mathbf{z}}$, where we have written the transverse part $\mathbf{K}$ of the wave vector $\mathbf{k}$ as $\mathbf{K} = k_x\hat{\mathbf{x}} + k_y\hat{\mathbf{y}}$ and $k_x^2 + k_y^2 + k_z^2 = k_0^2$ with $k_0 = \omega_0/c$. Here with $\sigma = +1$ and $\sigma = -1$ we denote left and right circular polarization, respectively. The helicity basis vectors $\hat{\mathbf{e}}_{1,2}(\mathbf{K})$ together with $\hat{\mathbf{k}} = \mathbf{k}/k_0$ itself, form a "local" Cartesian reference frame $\{\hat{\mathbf{e}}_1(\mathbf{K}), \hat{\mathbf{e}}_2(\mathbf{K}), \hat{\mathbf{k}}\}$ attached to the wave vector $\mathbf{k}$, whose expression can be obtained from the "global" reference frame attached to the central direction of propagation $\hat{\mathbf{z}}$ of the beam and denoted with $\{\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}\}$, via a rotation by the angle $\theta = \arccos(\hat{\mathbf{k}} \cdot \hat{\mathbf{z}})$ around the axis $\hat{\mathbf{n}} = \hat{\mathbf{z}} \times \hat{\mathbf{k}}/|\hat{\mathbf{z}} \times \hat{\mathbf{k}}|$ [27]. Explicitly, the potential vector in the angular spectrum representation reads as:

$$\mathbf{A}(\mathbf{r}) = \frac{1}{2\pi} \sum_{\sigma=\pm1} \int a_\sigma(\mathbf{K})\hat{\mathbf{h}}_\sigma(\mathbf{K}) e^{ik_z z} e^{i\mathbf{K}\cdot\mathbf{R}} \mathrm{d}^2K, \tag{8}$$

where $\mathrm{d}^2K = \mathrm{d}k_x\mathrm{d}k_y$ and $k_z = \sqrt{k_0^2 - k_x^2 - k_y^2} \geq 0$ for the case of forward homogeneous fields considered here [26]. Substituting this expression into Eq. (7) and using the relation $\mathrm{Re}\left\{iA_z^*(\mathbf{r} \times \mathbf{A})\right\} = i\left[z(\mathbf{A}^* \times \mathbf{A}) - \mathbf{r} \cdot (\mathbf{A}^* \times \mathbf{A})\hat{\mathbf{z}}\right]/2$ we obtain after a straightforward calculation:

$$\mathbf{J}^{\mathrm{surf}} = \frac{1}{2} \sum_{\sigma=\pm1} \sigma \int |a_\sigma(\mathbf{K})|^2 \left(\frac{k_0}{k_z}\hat{\mathbf{z}} - \frac{\mathbf{k}}{k_0} \right) \mathrm{d}^2K. \tag{9}$$

This explicit form for $\mathbf{J}^{\mathrm{surf}}$ is the first main result of this Letter: What does it represent? The transverse part of the term under the sign of integration is equal to $-1/2$ times the standard AM density in Fourier space, namely $\mathbf{K}/k_0$. Conversely, the longitudinal component is a completely new term that can be written as $\kappa^2/(2\sqrt{1-\kappa^2})$, where $\kappa \equiv |\mathbf{K}|/k_0$. This contribution is $O(\kappa^2)$ and therefore disappears in the standard paraxial limit where such terms are not retained [25]. In order to understand how this surface part affects the total AM per unit length we must add it to the standard spin AM density originating from the second term in Eq. (4) and equal to

$$\mathbf{J}^{\mathrm{spin}} = \sum_{\sigma=\pm1} \sigma \int |a_\sigma(\mathbf{K})|^2 \frac{\mathbf{k}}{k_0} \mathrm{d}^2K. \tag{10}$$

By comparing Eqs. (9) and (10), we immediately recognize that $\mathbf{J}^{\mathrm{surf}}$ has the structure of a spin angular momentum, as anticipated in the discussion after Eq. (7). Finally, the total spin AM is simply given by the sum $\mathbf{J}_{\mathrm{tot}}^{\mathrm{spin}} = \mathbf{J}^{\mathrm{spin}} + \mathbf{J}^{\mathrm{surf}}$:

$$\mathbf{J}_{\mathrm{tot}}^{\mathrm{spin}} = \frac{1}{2} \sum_{\sigma=\pm1} \sigma \int |a_\sigma(\mathbf{K})|^2 \left(\frac{\mathbf{k}}{k_0} + \frac{k_0}{k_z}\hat{\mathbf{z}} \right) \mathrm{d}^2K. \tag{11}$$

In the paraxial limit the terms between parenthesis in the equation above reduce to $2\hat{\mathbf{z}}$ and we recover the well known paraxial result $\mathbf{J}^{\rm spin}_{\rm tot} = \hat{\mathbf{z}} \sum_\sigma \sigma \int |a_\sigma(\mathbf{K})|^2 \mathrm{d}^2K$. If we denote with $\theta = \arccos(\hat{\mathbf{k}} \cdot \hat{\mathbf{z}})$ the angle formed by the generic wave vector $\mathbf{k}$ with the axis z, then the longitudinal contribution to Eq. (11) can be written as $\hat{\mathbf{z}}(\cos\theta + 1/\cos\theta)/2$. Interestingly, this term has the same form of the intrinsic AM with respect to the z-axis of a beam when observed in a reference frame tilted by the angle θ with respect to the direction of propagation [23, 28].

For the sake of definiteness, in the remainder we assume that the beam is circularly polarized along the direction $\hat{\mathbf{f}} = (1/\sqrt{2})(\hat{\mathbf{x}} + i\Sigma\hat{\mathbf{y}})$, where $\Sigma = \pm 1$ is the configuration-space helicity assigned in the (global) laboratory reference frame $\{\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}\}$ and not to be confused with the momentum-space helicity σ defined with respect to the (local) wave vector frame $\{\hat{\mathbf{e}}_1(\mathbf{K}), \hat{\mathbf{e}}_2(\mathbf{K}), \hat{\mathbf{k}}\}$. Therefore we can write the angular spectrum of the beam as $a_\sigma(\mathbf{K}) = \tilde{a}(\mathbf{K})\alpha_\sigma(\mathbf{K})$, where $\alpha_\sigma(\mathbf{K}) = \hat{\mathbf{f}} \cdot \hat{\mathbf{h}}_\sigma(\mathbf{K})$ is simply the projection of the global polarization of the beam onto the local helicity basis. From this definition it follows that $\sum_\sigma |\alpha_\sigma|^2 = (1 + \kappa_z^2)/2$ and $\sum_\sigma \sigma |\alpha_\sigma|^2 = \kappa_z \Sigma$, where we have defined $\kappa_x \hat{\mathbf{x}} + \kappa_y \hat{\mathbf{y}} + \kappa_z \hat{\mathbf{z}} = \hat{\boldsymbol{\kappa}} \equiv \mathbf{k}/k_0$. Thus, according to Eqs. (5-6) we can write

$$\mathbf{J}^{\rm surf} = \frac{\Sigma}{2} \int |\tilde{a}(\mathbf{K})|^2 \left(\hat{\mathbf{z}} - \kappa_z \hat{\boldsymbol{\kappa}}\right) \mathrm{d}^2K, \tag{12a}$$

$$\mathbf{J}^{\rm spin} = \Sigma \int |\tilde{a}(\mathbf{K})|^2 \, \kappa_z \hat{\boldsymbol{\kappa}} \, \mathrm{d}^2K, \tag{12b}$$

and

$$\frac{U}{\omega_0} = \frac{1}{2} \int |\tilde{a}(\mathbf{K})|^2 \left(1 + \kappa_z^2\right) \mathrm{d}^2K. \tag{13}$$

Since $\mathbf{J}^{\rm surf} \propto \Sigma$, then irrespective of the form of $\tilde{a}(\mathbf{K})$, $\overline{\mathbf{J}^{\rm surf}}$ drops to zero for a non-circularly polarized beam thus revealing the spin, as opposed to orbital, character of the SSAM. From Eqs. (12) we can straightforwardly calculate the projection of the spin and surface AM onto the central direction of proagation z of the beam:

$$\mathbf{J}^{\rm surf} \cdot \hat{\mathbf{z}} = \frac{\Sigma}{2} \int |\tilde{a}(\mathbf{K})|^2 (1 - \kappa_z^2) \, \mathrm{d}^2K, \tag{14a}$$

$$\mathbf{J}^{\rm spin} \cdot \hat{\mathbf{z}} = \Sigma \int |\tilde{a}(\mathbf{K})|^2 \kappa_z^2 \, \mathrm{d}^2K. \tag{14b}$$

From these expressions it follows that

$$\begin{aligned} \left(\mathbf{J}^{\rm spin} + \mathbf{J}^{\rm surf}\right) \cdot \hat{\mathbf{z}} &= \frac{\Sigma}{2} \int |\tilde{a}(\mathbf{K})|^2 (1 + \kappa_z^2) \, \mathrm{d}^2K \\ &= \Sigma \frac{U}{\omega_0}, \end{aligned} \tag{15}$$

where Eq. (13) has been used. Now, by using the definition Eq. (5), we obtain

$$\left(\overline{\mathbf{J}^{\rm spin}} + \overline{\mathbf{J}^{\rm surf}}\right) \cdot \hat{\mathbf{z}} = \Sigma. \tag{16}$$

Equation (16) shows that the role of the SSAM term is to ensure that the z-component of spin AM per unit length along the propagation axis z is *exactly* equal to Σ times the energy per unit length of the beam, the latter being completely arbitrary. This our second main result.

Examples.—In the remainder of this Letter, we calculate explicitly Eq. (12) for Laguerre-Gaussian and Bessel beams to investigate whether orbital AM affects $\overline{\mathbf{J}^{\rm surf}}$ or not. Consider first the case of a Laguerre-Gaussian beam of waist w_0, angular aperture $\theta_0 = 2/(k_0 w_0)$ and with azimuthal and radial mode indexes ℓ and p, respectively [29]. In this case, from Eqs. (12-13) we obtain

$$\overline{\mathbf{J}^{\rm surf}} = \Sigma \frac{\beta}{1 - \beta} \hat{\mathbf{z}}, \quad \overline{\mathbf{J}^{\rm spin}} = \Sigma \frac{1 - 2\beta}{1 - \beta} \hat{\mathbf{z}}, \tag{17}$$

where $\beta = (\theta_0/2)^2 (2p + |\ell| + 1)$. Although β depends on $|\ell|$ such a dependence is not a consequence of the orbital AM of the beam (this would lead to a dependence from the signed quantity ℓ), but it comes from the radial dependence of the beam profile. This result should be compared with the ones presented in Ref. [6]. Note that for a fundamental Gaussian beam $\ell, p \to 0$ and $\beta \to \theta_0^2/4$.

Next, we consider the case of a ℓth-order Bessel beam of the form $E(r, \phi, z) = J_\ell(r k_0 \sin\vartheta_0) \exp(i\ell\phi) \exp(i z k_0 \cos\vartheta_0)$, where ϑ_0 is the angular aperture of the characteristic Bessel cone. A straightforward calculation shows that Eqs. (17) are still valid providing that $\beta = \sin^2 \vartheta_0/2$. Here β is independent from the orbital AM value ℓ, thus supporting the claim that the SSAM has a spin nature. Note that for both Laguerre-Gaussian and Bessel *paraxial* beams with $\theta_0 = \vartheta_0 \ll 1$, one has $\overline{\mathbf{J}^{\rm surf}} \sim O(\theta_0^2)$, namely $\overline{\mathbf{J}^{\rm surf}}$ becomes negligible.



FIG. 1. SSAM for a circularly polarized fundamental Gaussian beam(green line) and Bessel beam (blue line).

The absolute value of the “surface helicity” $\Sigma^{\rm surf} \equiv |\overline{\mathbf{J}^{\rm surf}} \cdot \hat{\mathbf{z}}|$, is depicted in Fig. 1 for both the case of a fundamental Gaussian beam (green solid line) and of a Bessel beam (blue solid line) for $\vartheta_0 = \theta_0 \in [0, \pi/2]$. From this figure it appears clearly that in the paraxial domain Σ^{surf} approaches zero and its contribution can be neglected for $\theta_0 \ll 1$. This is in complete agreement with the fact that in the paraxial limit $k_z \simeq k_0$, $\mathbf{K} \simeq \mathbf{0}$ and from Eq. (9) it follows that $\mathbf{J}^{\text{surf}} \to \mathbf{0}$. This term, however, plays an important role in the non paraxial case, where, depending on the beam divergence, it can even become comparable with the standard helicity value $|\Sigma| = 1$ [25].

Conclusions.— By means of a careful analysis of the standard procedure leading to the separation in the spin and the orbital parts of the AM of a light beam, we have shown that a self-consistent derivation of this result should account for the fact that the AM per unit length, as opposed to the total AM, is the physical quantity actually measured in most experimental optical setups. This practically means that thoroughly all the steps of such derivation 3D integrations over the whole space must be replaced with 2D integrations over the plane xy at $z =$ const. As a result of this replacement we have found that a new surface spin-like part, the SSAM, must be added to the conventional spin and orbital terms. Further, we have demonstrated that the SSAM term guarantees that the z-component of spin AM per unit length along the propagation axis z equals the helicity $\Sigma(= \pm 1)$ times the energy per unit length of the beam, and that this result is universal, namely valid for any beam either paraxial or non paraxial. Finally, by calculating the SSAM for the cases of Laguerre-Gaussian and Bessel beams, we have shown that this novel term may give a contribution comparable to the conventional spin angular momentum. We believe that our results may have a relevant conceptual impact upon the very lively and timely research field about AM of light [31, 32].

We are grateful to Konstantin Bliokh for fruitful and stimulating discussions.

---

[*] andrea.aiello@mpl.mpg.de

[1] J. C. Maxwell, *A Treatise on Electricity and Magnetism*, Dover (1954).
[2] J. H. Poynting, Proc. R. Soc. Lond. A Ser. A **82**, 560567 (1909)
[3] R. A. Beth, Phys. Rev. **50**,115 (1936).
[4] C. G. Darwin, Proc. R. Soc. Lond. A **136**, 36 (1932).
[5] L. Allen, M. W. Beijersbergen, R. J. C. Spreeuw and J. P. Woerdman, Phys. Rev. A **45**, 8185 (1992).
[6] S. M. Barnett and L. Allen, Opt. Comm. **110**, 670 (1994).
[7] S. J. van Enk, and G. Nienhuis, Europhys. Lett., **25**, 497 (1994).
[8] S. J. van Enk, and G. Nienhuis, J. Mod. Opt., **41**, 963 (1994).
[9] S. M. Barnett, J. Opt. B: Quantum Semiclass. Opt. **4**, S7 (2002).
[10] M. V. Berry, J. Opt. A: Pure Appl. Opt., **11**, 094001 (2009).
[11] C.-F. Li, Phys. Rev. A, **80**, 063814 (2009).
[12] K. Y. Bliokh, M. A. Alonso, E. A. Ostrovskaya, and A. Aiello Phys. Rev. A **82**, 063825 (2010).
[13] S. M. Barnett, J. Mod. Opt., **57**, 1339 (2010).
[14] I. Bialynicki-Birula and Z. Bialynicki-Birula, J. Opt. **13**, 064014 (2011).
[15] O. Hosten and P. Kwiat, *Science* **319**, 787 (2008).
[16] A. Aiello and J. P. Woerdman, Opt. Lett. **33**, 1437 (2008).
[17] M. Onoda, S. Murakami and N. Nagaosa, Phys. Rev. Lett. **93**, 083901 (2004).
[18] K. Y. Bliokh and Y. P. Bliokh, Phys. Rev. Lett. **96**, 073903 (2006).
[19] A. Y. Bekshaev, J. Opt. A: Pure Appl. Opt. **11**, 094003 (2009).
[20] A. Aiello, N. Lindlein, C. Marquardt and G. Leuchs, *Phys. Rev. Lett.* **103**, 100401 (2009)
[21] S. Franke-Arnold, L. Allen and M. Padgett, *Laser & Photon. Rev.* **2**, 299 (2008).
[22] *Optical Angular Momentum*, edited by L. Allen, S. M. Barnett and M. J. Padgett (Institute of Physics, Bristol, England, 2003).
[23] K. Y. Bliokh, A. Aiello and M. Alonso in *The Angular Momentum of Light*, edited by D. L. Andrews and M. Babiker, Cambridge University Press (2012).
[24] J. H. Crichton and P. L. Marston, Electronic Journal of Differential Equations, http://www.emis.de/journals/EJDE/, pp.37-50 (2000).
[25] H. A. Haus and J. L. Pan, Am. J. Phys. **61**, 818 (1993).
[26] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics*, Cambridge University Press (1995).
[27] A. Aiello, C. Marquardt and G. Leuchs, Phys. Rev. A **81**, 053838 (2010).
[28] We acknowledge Konstantin Bliokh for having pointed out this analogy.
[29] A. E. Siegman, *Lasers*, (University Science Books, USA, 1986).
[30] A. Aiello and J. P. Woerdman, Opt. Lett. **36**, 543 (2011).
[31] O. G. Rodríguez-Herrera, D. Lara, K. Y. Bliokh, E. A. Ostrovskaya, and C. Dainty, Phys. Rev. Lett. **104**, 253601 (2010).
[32] P. Banzer, U. Peschel, S. Quabis, G. Leuchs, Opt. Expr. **18**, 10905 (2010).